

READ SAFETY DIRECTIONS BEFORE OPENING OR USING. FOR ANIMAL TREATMENT ONLY.



VETRAZIN[®]

LIQUID

Sheep Blowfly Treatment

Active Constituent: **500 g/L CYROMAZINE**

READ DIRECTIONS FOR USE BEFORE USING THIS PRODUCT

PHARM SMART AUSTRALIA PTY LTD.

Suite 11, Level 1, 240 Waterworks Road, Ashgrove QLD 4060, Australia

7 Litres



Protects long wool sheep for up to 14 weeks from strike by blowfly (*Lucilia cuprina*), including organophosphate resistant strains, when applied by jetting or dipping.

Pharmsmart

10489

VETRAZIN

LIQUID

DOSAGE AND ADMINISTRATION

Animal/Pest	Treatment	Initial Charge	Topping Up	Critical Comments
Sheep Blowfly Control (<i>Lucilia cuprina</i>)	Jetting	2 L per 1000 L	—	<ol style="list-style-type: none">1. Apply 2 to 4 L of jetting fluid per sheep, depending on sheep size and wool length.2. Ensure saturation to the skin.3. Pay particular attention to treatment of the crutch, pizzlies and polls.4. If used in jetting races a reduced period of protection may result.5. Heavy rain following the application of this product could diminish the period of protection.
	Dipping (Plunge, Shower and Constant Replenishment)	2 L per 1000 L	2 L per 1000 L	<ol style="list-style-type: none">1. Dip sheep at onset of fly wave.2. Thoroughly saturate sheep. When using a shower dip, run top and bottom sprays independently so as to ensure maximum pressure and therefore penetration.3. The product is a non-stripping dip so there is no need to reinforce during dipping.4. Dipping sheep heavily infested with grass seeds is not recommended.5. Heavy rain following the application of this product could diminish the period of protection.
	Dressing	2 L per 1000 L	—	<ol style="list-style-type: none">1. Clip the wool surrounding the strike, treat the struck area and the surrounding wool with a recommended knockdown product then thoroughly jet the sheep with Vetrazin.2. This product is a slow killing larvicide and when used on its own, larvae may take 3 to 4 days to die.

DIRECTIONS FOR USE

Restraints

DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.

DO NOT USE on sheep with less than 6 weeks wool.

Contraindications

Should not be used on open wounds, e.g. those resulting from marking or mulesing operations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

MILK: DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.

WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1800 979 692 or the APVMA website (www.apvma.gov.au/residues). Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.