

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**



VETRAZIN[®]

SPRAY-ON

Sheep Blowfly Treatment

Active Constituent: 60g/L CYROMAZINE

**READ DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT**

PHARM SMART AUSTRALIA PTY LTD.
Suite 11, Level 1
240 Waterworks Road
Ashgrove QLD 4060
Australia

25 Litres
10487



When applied as directed will protect long wool sheep against fly strike for 11 weeks.

Pharmsmart

VETRAZIN[®]

SPRAY-ON

DIRECTIONS FOR USE

Restraints

- DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.
- DO NOT USE on sheep with less than 6 weeks wool.

Contraindications

Should not be used on open wounds, e.g. those resulting from marking or mulesing operations.

DOSAGE AND ADMINISTRATION

FOR THE PREVENTION OF BODY STRIKE

Months since shearing	Total volume/sheep (mL)	Volume per treatment band (mL)	Number of bands	Number of sheep treated per 25 L
up to 6	34	17	2	735
7	38	19	2	657
8	44	22	2	568
9	50	25	2	500
10	56	28	2	446
Lambs less than 15 kg	17	17	1	1470

FOR THE PREVENTION OF BODY & CRUTCH STRIKE

Months since shearing	Total volume/sheep (mL)	Volume per treatment band (mL)	Number of bands	Number of sheep treated per 25 L
up to 6	51	17	3	490
7	57	19	3	439
8	66	22	3	378
9	75	25	3	333
10	84	28	3	297
Lambs less than 15 kg	34	17	2	735

FOR THE PREVENTION OF POLL STRIKE

Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of Vetrazin Spray-On to the areas requiring protection, around the poll and horns.

CRITICAL COMMENTS

- Use only a Pharmsmart recommended Spray-On applicator gun.
- Ensure the applicator gun is clean.
- Calibrate applicator gun using an accurate volume measure.
- Ensure all areas to be protected are treated.
- When preventing body strike ensure bands overlap along the midline - see diagram.
- This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die.
- Should not be mixed with any other product.
- Should not be applied during gusty or windy weather as spray drift will reduce the period of protection.
- Heavy rain following the application of this product could diminish the period of protection.
- Vetrazin Spray-On is coloured for band identification. This scourable dye will disappear shortly after treatment.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption

MILK: DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption


WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than 28 days before slaughter for export.

The ESI on this label was correct at the time of label approval.

Before using this product, confirm the current ESI from the manufacturer on 1800 979 692 or the APVMA website (www.apvma.gov.au/residues). Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.



GENERAL INSTRUCTIONS

DO NOT DILUTE. Apply directly from the container using a Pharmsmart recommended Spray-On applicator gun. Except for small lambs under 15 kg, the dose volume is determined by the number of months since shearing. Apply Vetrazin Spray-On according to the diagrams and tables in the Directions for Use.

Vetrazin Spray-On may be applied to all breeds of sheep, providing protection against fly strike. Vetrazin Spray-On is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25 cm above the fleece, to achieve a 15 cm wide band at each pass.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Ph: 13 1126).

ADDITIONAL USER SAFETY

Sheep Rehhandling Interval: After treatment wait until sheep are dry before rehhandling.

Protection of Wildlife, Fish, Crustacea and Environment

DO NOT contaminate dams, rivers or streams with the product or used containers.

Disposal

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

Storage

Store below 30°C (Room Temperature). Store in original container tightly closed in a dry, cool place. Do not store in direct sunlight. Do not re-use the container. Keep out of reach of children.

May cause an allergic skin reaction. Causes serious eye irritation. Toxic to aquatic life with long lasting effects. If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone 1800 979 692.

WARRANTY AND EXCLUSION OF LIABILITY

This product is warranted fit for the purposes specifically recommended by Pharm Smart Australia Pty Ltd when used strictly as directed on this label. All other warranties and obligations or liabilities, whether expressed or implied by statute or otherwise, are excluded to the full extent that exclusion is permitted by law.

CUSTOMER INFORMATION Australia: 1800 979 692 TOLL FREE

© Vetrazin is a trademark owned or licensed by Pharm Smart Australia Pty Ltd, its subsidiaries or affiliates. APVMA Approval Number: 39979/0614

Batch Number:	
Expiry Date:	

