

DANGER
FOR ANIMAL TREATMENT ONLY
SHAKE WELL BEFORE USE



VETRAZIN[®]

SPRAY-ON

**For the preventative control
of blowfly strike on sheep**

ACTIVE CONSTITUENT:
60 g/L Cyromazine

Contains 16.7 g/L
glacial acetic acid



NET CONTENT:
25 Litres

Pharmsmart

DANGER
KEEP OUT OF REACH OF CHILDREN
 Dangerous to the environment
 Corrosive



PLEASE READ THE LABEL COMPLETELY BEFORE USE

Safety Directions: May cause skin irritation. Avoid skin contact. May cause eye damage. Avoid contact with eyes. Wear eye and face protection. Cyromazine possibly may affect blood and the haemopoietic system. Acetic acid possibly may damage lungs following inhalation. Do not breathe fumes, vapours, or spray. Wash hands thoroughly after handling. Avoid release to the environment. Wear protective gloves, clothing, and eye protection. Handle with care.

First Aid: If medical advice is needed, have product container or label at hand. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/ attention. Take off contaminated clothing and wash before re-use. IF exposed or concerned: Get medical advice/ attention. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Contact the National Poisons Centre **0800 POISON (0800 764 766)** or a doctor immediately.

Environmental Protection: Harmful to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container.

Emergency response: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely. If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to www.pharmsmart.co.nz.

Disposal: Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse container with water and dispose of rinsate away from waterways. Crush or puncture and bury in a suitable landfill. Do NOT use container for any other purpose.

*For further details see the Safety Data Sheet.

VETRAZIN[®]

SPRAY-ON

DIRECTIONS FOR USE


Apply VETRAZIN SPRAY-ON through a suitable clean applicator gun. The product should be sprayed on in bands with the gun held 20 cm above the wool during application. VETRAZIN SPRAY-ON should be applied onto those areas where maximum protection against flystrike is required. VETRAZIN SPRAY-ON applied between the rump and shoulders should be sprayed on in even bands as shown (see diagram). The product sprayed onto the rump should be applied in an arc over the area shown (see diagram). The dose rate and number of bands to be applied are shown in the table. Note: The dose volume applied should be based on the weight of the heaviest sheep in the mob.

Application and dose rate of VETRAZIN SPRAY-ON for the protection of all the commonly struck areas of the sheep.


Live weight (kg)	11-15		16-20		21-30		31-40		41-50		51-60		61-70	
	2	2	3	3	4	4	5							
Number of bands	2	2	3	3	4	4	5							
Volume per band (mL)	7.5	10	10	13.3	12.5	15	14							
Total volume applied (mL)	15	20	30	40	50	60	70							
Sheep treated per 10 L	666	500	333	250	200	166	142							

Application and dose rate of VETRAZIN SPRAY-ON at docking for the prevention of fly strike.


Apply between 5 mL and 10 mL over the crutch ensuring that sufficient product to cover the area to be protected is applied (see diagram).




2 bands
 Sheep between 11-20 kg



3 bands
 Sheep between 21-40 kg



4 bands
 Sheep between 41-60 kg



5 bands
 Sheep between 61-70 kg

STORAGE

Store in original container, tightly closed, away from direct sunlight, in a cool, dry place, avoiding temperatures below 0 °C and above 35 °C.

WARRANTY

The manufacturer of this animal remedy extends/grants to the purchaser a warranty that this animal remedy is reasonably fit for the purposes for which its use is recommended, provided that the purchaser uses the remedy only for the purposes for which it is recommended and strictly in accordance with the directions on this container.

Registered Pursuant to the ACVM Act 1997, No. A005717
 See www.foodsafety.govt.nz for registration conditions.

Approved Pursuant to the HSNO Act 1996, Approval Code: HSR100759
 See www.epa.govt.nz for approval conditions.

Registered to: Pharmsmart NZ Ltd,
 Level 8, 430 Victoria Street,
 Hamilton 3204,
 New Zealand
 HELPLINE: Phone 0800 694 401
www.pharmsmart.co.nz

NET CONTENT 25 LITRES

®VETRAZIN is a registered trademark of Pharmsmart and its affiliates.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

WITHHOLDING PERIODS

- MEAT:** Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of cessation of the last treatment.
- MILK:** Milk intended for sale for human consumption must be discarded during treatment for not less than 35 days following the last treatment.

WOOL RESIDUES

Ensure the time between dipping and subsequent shearing is at least 2 months, except in the case of spot treatment of flystrike. For advice on your dipping programme, contact your supplier.

ACTIVE CONSTITUENT

VETRAZIN SPRAY-ON contains 60 g/L Cyromazine in the form of an aqueous concentrate.

GENERAL INFORMATION

VETRAZIN SPRAY-ON is an easy to use spray-on, which will give up to 6 weeks residual protection against flystrike caused by all the commonly found strike flies including the Australian blowfly (*Lucilia cuprina*). VETRAZIN SPRAY-ON can be used on all breeds of sheep. VETRAZIN SPRAY-ON is suitable for use on sheep between 4 and 12 weeks after shearing. A shorter period of protection may occur if VETRAZIN SPRAY-ON is applied incorrectly, if there is excessive fleece soiling or if wool infections such as mycotic dermatitis (fleece rot) are present. Prolonged heavy rain following treatment may also reduce the protection period provided by VETRAZIN SPRAY-ON.

Batch No:	
Expiry:	

Pharmsmart



9 359003 000052 10503